

\$

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 18:53:42 ON 20 JUN 2009

=> b ca

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.22	0.22

FILE 'CA' ENTERED AT 18:54:00 ON 20 JUN 2009

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 18 Jun 2009 VOL 150 ISS 26

FILE LAST UPDATED: 18 Jun 2009 (20090618/ED)

REVISED CLASS FIELDS (/NCL) LAST RELOADED: Arp 2009

USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Feb 2009

CA now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2008.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s apm1 or adiponectin

173 APM1

5472 ADIPONECTIN

L1 5556 APM1 OR ADIPONECTIN

=> s l1 and polyclonal antibod?

40047 POLYCLONAL

542111 ANTIBOD?

20073 POLYCLONAL ANTIBOD?

(POLYCLONAL(W)ANTIBOD?)

L2 15 L1 AND POLYCLONAL ANTIBOD?

=> s l2 and native(w)(apm1 or adiponectin)

156938 NATIVE

173 APM1

5472 ADIPONECTIN  
 2 NATIVE(W) (APM1 OR ADIPONECTIN)  
 L3 0 L2 AND NATIVE(W) (APM1 OR ADIPONECTIN)

=> s l2 and native  
 156938 NATIVE  
 L4 0 L2 AND NATIVE

=> s l2 and latex  
 75640 LATEX  
 L5 2 L2 AND LATEX

=> d ti ab 1-2

L5 ANSWER 1 OF 2 CA COPYRIGHT 2009 ACS on STN  
 TI Determination of \*\*\*adiponectin\*\*\* in serum using a \*\*\*latex\*\*\*  
 particle-enhanced turbidimetric immunoassay with an automated analyzer  
 AB \*\*\*Adiponectin\*\*\* is an adipose-derived hormone that plays a role in  
 regulating metabolic processes such as fat partitioning and lipid and  
 glucose metab. Quantification of \*\*\*adiponectin\*\*\* is useful for  
 obtaining information on metabolic syndrome, but there is no rapid method  
 to measure \*\*\*adiponectin\*\*\* for clin. use. The authors developed a  
 rapid and sensitive \*\*\*latex\*\*\* particle-enhanced turbidimetric  
 immunoassay (LTIA) using a \*\*\*latex\*\*\* bead-immobilized anti-  
 \*\*\*adiponectin\*\*\* \*\*\*polyclonal\*\*\* \*\*\*antibody\*\*\*. The assay  
 was performed on a Hitachi H7170 analyzer and evaluated for validity as a  
 method to quantitate \*\*\*adiponectin\*\*\*, in parallel with the ELISA.  
 Diln. tests using LTIA showed linearity from 0.25 to 30 .mu.g/mL.  
 Within-run CV and total CV were obtained in the range 0.8-1.9% and  
 1.1-2.0%, resp. No interference was obsd. in the testing of specimens  
 contg. potentially interfering substances such as bilirubin,  
 ditaurobilirubin, Hb triglyceride, rheumatoid factor, type IV collagen,  
 fibronectin, and complement factor (Clq). A strong correlation between  
 LTIA and ELISA was confirmed (n = 30, r = 0.990, y = 0.95x + 0.39). The  
 LTIA assay is applicable to quantitating the serum concn. of  
 \*\*\*adiponectin\*\*\*. This assay is more convenient and faster than ELISA  
 and suitable for clin. routine anal.

L5 ANSWER 2 OF 2 CA COPYRIGHT 2009 ACS on STN  
 TI \*\*\*Latex\*\*\* reagent for \*\*\*adiponectin\*\*\* analysis, and  
 \*\*\*adiponectin\*\*\* analysis method  
 AB A \*\*\*latex\*\*\* reagent for \*\*\*adiponectin\*\*\* anal. is provided,  
 which comprises a suspension of \*\*\*latex\*\*\* particles carrying a  
 substance (e.g., anti- \*\*\*adiponectin\*\*\* \*\*\*polyclonal\*\*\*  
 \*\*\*antibody\*\*\*) capable of specifically binding with  
 \*\*\*adiponectin\*\*\*  
 . Also provided is a method for \*\*\*adiponectin\*\*\* anal., which  
 comprises: (1) a step for obtaining a biol. liq. potentially contg.  
 \*\*\*adiponectin\*\*\*; and (2) a step for contacting the biol. liq.  
 obtained  
 in the step (1) as it is with a suspension of \*\*\*latex\*\*\* particles  
 carrying a substance capable of specifically binding with  
 \*\*\*adiponectin\*\*\*, and optically analyzing the resultant mixt. to det.  
 the degree of agglutination of the \*\*\*latex\*\*\* particles. According  
 to this \*\*\*latex\*\*\* reagent for \*\*\*adiponectin\*\*\* anal. and this  
 \*\*\*adiponectin\*\*\* anal. method, it is not required to dil. or pretreat

biol. liq. as a test sample beforehand, and the anal. is rapidly and conveniently performed without limiting a measurement facility.

=> d all 1-2

L5 ANSWER 1 OF 2 CA COPYRIGHT 2009 ACS on STN

AN 145:391809 CA <<LOGINID::20090620>>

ED Entered STN: 02 Nov 2006

TI Determination of \*\*\*adiponectin\*\*\* in serum using a \*\*\*latex\*\*\* particle-enhanced turbidimetric immunoassay with an automated analyzer

AU Nishimura, Ayako; Sawai, Tokio

CS Division of Research and Development, Mitsubishi Kagaku Iatron Inc., 1460-6 Mitodai, Tako-machi, Katori-gun, Chiba-ken, 289-2247, Japan

SO Clinica Chimica Acta (2006), 371(1-2), 163-168

CODEN: CCATAR; ISSN: 0009-8981

PB Elsevier Ltd.

DT Journal

LA English

CC 9-10 (Biochemical Methods)

AB \*\*\*Adiponectin\*\*\* is an adipose-derived hormone that plays a role in regulating metabolic processes such as fat partitioning and lipid and glucose metab. Quantification of \*\*\*adiponectin\*\*\* is useful for obtaining information on metabolic syndrome, but there is no rapid method to measure \*\*\*adiponectin\*\*\* for clin. use. The authors developed a rapid and sensitive \*\*\*latex\*\*\* particle-enhanced turbidimetric immunoassay (LTIA) using a \*\*\*latex\*\*\* bead-immobilized anti-\*\*\*adiponectin\*\*\* \*\*\*polyclonal\*\*\* \*\*\*antibody\*\*\*. The assay was performed on a Hitachi H7170 analyzer and evaluated for validity as a method to quantitate \*\*\*adiponectin\*\*\*, in parallel with the ELISA. Diln. tests using LTIA showed linearity from 0.25 to 30 .mu.g/mL. Within-run CV and total CV were obtained in the range 0.8-1.9% and 1.1-2.0%, resp. No interference was obsd. in the testing of specimens contg. potentially interfering substances such as bilirubin, ditauobilirubin, Hb triglyceride, rheumatoid factor, type IV collagen, fibronectin, and complement factor (Clq). A strong correlation between LTIA and ELISA was confirmed (n = 30, r = 0.990, y = 0.95x + 0.39). The LTIA assay is applicable to quantitating the serum concn. of \*\*\*adiponectin\*\*\*. This assay is more convenient and faster than ELISA and suitable for clin. routine anal.

ST \*\*\*adiponectin\*\*\* detn serum \*\*\*latex\*\*\* particle turbidimetric immunoassay automated analyzer; metabolic syndrome \*\*\*adiponectin\*\*\* serum immunoturbidimetry automatic analyzer

IT Cytokines

RL: ANT (Analyte); ANST (Analytical study)

( \*\*\*adiponectin\*\*\* ; detn. of \*\*\*adiponectin\*\*\* in serum using \*\*\*latex\*\*\* particle-enhanced turbidimetric immunoassay with automated analyzer)

IT Blood analysis

Human

Immunoturbidimetry

Metabolic disorders

(detn. of \*\*\*adiponectin\*\*\* in serum using \*\*\*latex\*\*\* particle-enhanced turbidimetric immunoassay with automated analyzer)

RE.CNT 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

(1) Anon; JAMA 2001, V285, P2486

- (2) Arita, Y; Biochem Biophys Res Commun 1999, V257, P79 CA
- (3) Arita, Y; Circulation 2002, V105, P2893 CA
- (4) Bland, J; Lancet 1986, V1, P307 MEDLINE
- (5) Daimon, M; Diabetes Care 2003, V26, P2015 CA
- (6) Haluzik, M; Physiol Res 2004, V53, P123 CA
- (7) Hansa, T; Clin Chem 1997, V43, P109
- (8) Hotta, K; Arterioscler Thromb Vasc Med 2000, V20, P1595 CA
- (9) Hotta, K; Diabetes 2001, V50, P1126 CA
- (10) Kumada, M; Arterioscler Thromb Vasc Biol 2003, V23, P85 CA
- (11) Lindsay, R; Lancet 2002, V360, P57 CA
- (12) Lowry, O; J Biol Chem 1951, V193, P265 CA
- (13) Maeda, K; Biochem Biophys Res Commun 1996, V221, P286 CA
- (14) Maeda, K; Gene 1997, V190, P227 CA
- (15) Okamoto, Y; Circulation 2002, V106, P2767 CA
- (16) Okamoto, Y; Horm Metab Res 2000, V32, P47 CA
- (17) Ouchi, N; Circulation 1999, V100, P2473 CA
- (18) Ouchi, N; Circulation 2000, V102, P1296 CA
- (19) Ouchi, N; Circulation 2001, V103, P1057 CA
- (20) Ouchi, N; Hypertension 2003, V42, P231 CA
- (21) Pauli, S; Clin Chem 2004, V50, P219
- (22) Pischon, T; JAMA 2004, V291, P1730 CA
- (23) Ryo, M; Circ J 2004, V68, P975 CA
- (24) Tsukinoki, R; Lipids Health Dis 2005, V4, P27

L5 ANSWER 2 OF 2 CA COPYRIGHT 2009 ACS on STN

AN 141:310247 CA <<LOGINID::20090620>>

ED Entered STN: 28 Oct 2004

TI \*\*\*Latex\*\*\* reagent for \*\*\*adiponectin\*\*\* analysis, and  
 \*\*\*adiponectin\*\*\* analysis method

IN Tachikawa, Tetsuya; Akamatsu, Suguru; Sawai, Tokio; Nishimura, Fumiko  
 PA Mitsubishi Kagaku Iatron, Inc., Japan; Otsuka Pharmaceutical Co., Ltd.  
 SO PCT Int. Appl., 26 pp.

CODEN: PIXXD2

DT Patent

LA Japanese

IC ICM G01N033-53

ICS G01N033-543

CC 9-10 (Biochemical Methods)

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 2004086040	A1	20041007	WO 2004-JP4083	20040324
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	AU 2004223553	A1	20041007	AU 2004-223553	20040324
	AU 2004223553	B2	20081204		
	CA 2520438	A1	20041007	CA 2004-2520438	20040324
	EP 1607742	A1	20051221	EP 2004-723044	20040324

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK  
 US 20070037207 A1 20070215 US 2005-550324 20050923  
 PRAI JP 2003-80763 A 20030324  
 WO 2004-JP4083 W 20040324

# CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 2004086040	ICM	G01N033-53
	ICS	G01N033-543
	IPCI	G01N0033-53 [ICM,7]; G01N0033-543 [ICS,7]
	IPCR	G01N0033-53 [I,C*]; G01N0033-53 [I,A]; G01N0033-543 [I,C*]; G01N0033-543 [I,A]; G01N0033-68 [I,C*]; G01N0033-68 [I,A]
AU 2004223553	ECLA	G01N033/543D; G01N033/68V
	IPCI	G01N0033-53 [I,C*]; G01N0033-53 [I,A]; G01N0033-543 [I,C*]; G01N0033-543 [I,A]; G01N0033-68 [I,C*]; G01N0033-68 [I,A]
	IPCR	G01N0033-53 [I,C*]; G01N0033-53 [I,A]; G01N0033-543 [I,C*]; G01N0033-543 [I,A]; G01N0033-68 [I,C*]; G01N0033-68 [I,A]
CA 2520438	ECLA	G01N033/543D; G01N033/68V
	IPCI	G01N0033-53 [ICM,7]; G01N0033-543 [ICS,7]
	IPCR	G01N0033-53 [I,C*]; G01N0033-53 [I,A]; G01N0033-543 [I,C*]; G01N0033-543 [I,A]; G01N0033-68 [I,C*]; G01N0033-68 [I,A]
EP 1607742	ECLA	G01N033/543D; G01N033/68V
	IPCI	G01N0033-53 [ICM,7]; G01N0033-543 [ICS,7]
	IPCR	G01N0033-53 [I,C*]; G01N0033-53 [I,A]; G01N0033-543 [I,C*]; G01N0033-543 [I,A]; G01N0033-68 [I,C*]; G01N0033-68 [I,A]
US 20070037207	ECLA	G01N033/543D; G01N033/68V
	IPCI	G01N0033-53 [I,A]; G01N0033-551 [I,A]
	NCL	435/007.100; 436/524.000
AB	A	***latex*** reagent for ***adiponectin*** anal. is provided, which comprises a suspension of ***latex*** particles carrying a substance (e.g., anti- ***adiponectin*** ***polyclonal*** ***antibody*** ) capable of specifically binding with ***adiponectin*** . Also provided is a method for ***adiponectin*** anal., which comprises: (1) a step for obtaining a biol. liq. potentially contg. ***adiponectin*** ; and (2) a step for contacting the biol. liq. obtained in the step (1) as it is with a suspension of ***latex*** particles carrying a substance capable of specifically binding with ***adiponectin*** , and optically analyzing the resultant mixt. to det. the degree of agglutination of the ***latex*** particles. According to this ***latex*** reagent for ***adiponectin*** anal. and this ***adiponectin*** anal. method, it is not required to dil. or pretreat a biol. liq. as a test sample beforehand, and the anal. is rapidly and conveniently performed without limiting a measurement facility.
ST		***adiponectin*** analysis ***latex*** agglutination reagent antibody
IT		Antibodies and Immunoglobulins
		RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses) ( ***adiponectin*** anal. method using ***latex*** agglutination

```

immunoassay reagent)
IT   Cytokines
      RL: ANT (Analyte); ANST (Analytical study)
      (   ***adiponectin***   ;   ***adiponectin***   anal. method using
        ***latex***   agglutination immunoassay reagent)
IT   Agglutination test
      (   ***latex***   ;   ***adiponectin***   anal. method using
        ***latex***   agglutination immunoassay reagent)
IT   ***Latex***
      (particles;   ***adiponectin***   anal. method using   ***latex***
        agglutination immunoassay reagent)

```

RE.CNT 4        THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD  
RE

- (1) A&t Corp; JP 20014624 A 2001
- (2) Otsuka Pharmaceutical Co Ltd; EP 1033134 A 1999 CA
- (3) Otsuka Pharmaceutical Co Ltd; US 6461821 B 1999 CA
- (4) Otsuka Pharmaceutical Co Ltd; WO 9921577 A 1999 CA

=> d his

(FILE 'HOME' ENTERED AT 18:53:42 ON 20 JUN 2009)

FILE 'CA' ENTERED AT 18:54:00 ON 20 JUN 2009

```

L1      5556 S APM1 OR ADIPONECTIN
L2      15 S L1 AND POLYCLONAL ANTIBOD?
L3      0 S L2 AND NATIVE(W) (APM1 OR ADIPONECTIN)
L4      0 S L2 AND NATIVE
L5      2 S L2 AND LATEX

```

=> logoff y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	31.85	32.07
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-3.12	-3.12

STN INTERNATIONAL LOGOFF AT 18:56:58 ON 20 JUN 2009